THE MORRIS LAW FIRM, P.C.

INTELECTUAL PROPERTY LAW INCLUDING PATENT, TRADEMARK COPYRIGHT LAW, UNFAIR COMPETITION AND RELATED MATTERS 10777 WESTHEIMER, SUITE 1100

HOUSTON, TEXAS 77042

RECEIVED CENTRAL FAX CENTER

<u>MAULING ADDRESS:</u> P.O. BOX 420787 HOUSTON, TEXAS 77242-0787

CUSTOMER No. 23770

JUN 05 2009

Certificate of Facsimile Transmission Under 37 CFR 1.8

I hereby certify that this correspondence is being transmitted via facsimile transmission in accordance with 37 CFR 1.8 on the date indicated below addressed to:

Commissioner for Patents PO BOX 1450 Alexandria, VA 22313-1450

on 06-05-09 (571) 273-8300 No. of Pages including this cover sheet: 5

| Date | Facsimile Number | Typed or Printed Name | Typ

Attached are the following pages:

 Omitted Attachment to Response to Office Action Mailed March 24, 2009 (erroneously FAXed to Mail Stop AF) [4 pages]

Applicant: _	GUENTHER, et al.	Group Art Unit:	1797
Serial No.: _	10/521,700	Examiner:	Lessanework T. Seifu
Filing Date: _	01/19/2005	Atty. Docket No.:	TS8579US

Title: "Process to Generate Heat"

Page 1 of 4


Evaporation

From Wikipedia, the free encyclopedia

Evaporation is the slow vaporization of a liquid and the reverse of condensation. A type of phase transition, it is the process by which molecules in a liquid state (e.g. water) spontaneously become gaseous (e.g. water vapor). Generally, evaporation can be seen by the gradual disappearance of a liquid from a substance when exposed to a significant volume of gas. Vaporization and evaporation however, are not entirely the same processes. For example, substances like caesium, francium, gallium, bromine, rubidium and mercury may vaporize, but they do not evaporate as such.

On average, the molecules in a glass of water do not have enough heat energy to escape from the liquid, or else the liquid would turn into vapor quickly (see boil). When the molecules collide, they transfer energy to each other in varying degrees, based on how they collide. Sometimes the transfer is so one-sided for a molecule near the surface that it ends up with enough energy to escape.

Liquids that do not evaporate visibly at a given temperature in a given gas (e.g. cooking oil at room temperature) have molecules that do not tend to transfer energy to each other in a pattern sufficient to frequently give a molecule the heat energy necessary to turn into vapor. However, these liquids are evaporating, it's just that the process is much slower and thus significantly less visible.



(713) 334-5157

Water vapor that has evaporated and disappeared from hot tea condenses into visible droplets. Gaseous water is invisible, but the clouds of water droplets are evidence of evaporation followed by condensation. It is in the water cycle

Evaporation is an essential part of the water cycle. Solar energy drives evaporation of water from oceans, lakes, moisture in the soil, and other sources of water. In hydrology, evaporation and transpiration (which involves evaporation within plant stomata) are collectively termed evapotranspiration. Evaporation is caused when water is exposed to air and the liquid molecules turn into water vapor which rises up and forms clouds.

Contents

- 1 Theory
 - 1.1 Evaporative equilibrium
- 2 Factors influencing the rate of evaporation
- 3 Applications
 - 3.1 Combustion vaporization
 - 3.2 Film deposition
- 4 See also
- 5 References
- 6 External links

Theory

See also: Kinetic theory

Page 2 of 4


For molecules of a liquid to evaporate, they must be located near the surface, be moving in the proper direction, and have sufficient kinetic energy to overcome liquid-phase intermolecular forces.^[1] Only a small proportion of the molecules meet these criteria, so the rate of evaporation is limited. Since the kinetic energy of a molecule is proportional to its temperature, evaporation proceeds more quickly at higher temperatures. As the faster-moving molecules escape, the remaining molecules have lower average kinetic energy, and the temperature of the liquid thus decreases. This phenomenon is also called evaporative cooling. This is why evaporating sweat cools the human body. Evaporation also tends to proceed more quickly with higher flow rates between the gaseous and liquid phase and in liquids with higher vapor pressure. For example, laundry on a clothes line will dry (by evaporation) more rapidly on a windy day than on a still day. Three key parts to evaporation are heat, humidity and air movement.

On a molecular level, there is no strict boundary between the liquid state and the vapor state. Instead, there is a Knudsen layer, where the phase is undetermined. Because this layer is only a few molecules thick, at a macroscopic scale a clear phase transition interface can be seen.

Evaporative equilibrium

If evaporation takes place in a closed vessel, the escaping molecules accumulate as a vapor above the liquid. Many of the molecules return to the liquid, with returning molecules becoming more frequent as the density and pressure of the vapor increases. When the process of escape and return reaches an equilibrium, [1] the vapor is said to be "saturated," and no further change in either vapor pressure and density or liquid temperature will occur. For a system consisting of vapor and liquid of a pure substance, this equilibrium state is directly related to the vapor pressure of the substance, as given by the Clausius-Clapeyron relation:

$$\ln\left(\frac{P_2}{P_1}\right) = -\frac{\Delta H_{vap}}{R} \left(\frac{1}{T_2} - \frac{1}{T_1}\right)$$



where P_1 , P_2 are the vapor pressures at temperatures T_1 , T_2 respectively, $\Delta H_{\rm vap}$ is the enthalpy of vaporization, and R is the universal gas constant. The rate of evaporation in an open system is related to the vapor pressure found in a closed system. If a liquid is heated, when the vapor pressure reaches the ambient pressure the liquid will boil.

The ability for a molecule of a liquid to evaporate is largely based on the amount of kinetic energy an individual particle may possess. Even at lower temperatures, individual molecules of a liquid can evaporate if they have more than the minimum amount of kinetic energy required for vaporization.

But vaporization is not only the process of a change of state from liquid to gas but it is also a change of state from a solid to gas. This process is also known as sublimation but can also be known as vaporization.

Factors influencing the rate of evaporation

Page 3 of 4

Concentration of the substance evaporating in the air

If the air already has a high concentration of the substance evaporating, then the given substance will evaporate more slowly.

Concentration of other substances in the air

If the air is already saturated with other substances, it can have a lower capacity for the substance evaporating.

Concentration of other substances in the liquid (impurities)

If the liquid contains other substances, it will have a lower capacity for evaporation.

Flow rate of air

Jun 05 2009 18:20

This is in part related to the concentration points above. If fresh air is moving over the substance all the time, then the concentration of the substance in the air is less likely to go up with time, thus encouraging faster evaporation. This is the result of the boundary layer at the evaporation surface decreasing with flow velocity, decreasing the diffusion distance in the stagnant layer.

Inter-molecular forces

The stronger the forces keeping the molecules together in the liquid state, the more energy one must get to escape.

Pressure

In an area of less pressure, evaporation happens faster because there is less exertion on the surface keeping the molecules from launching themselves.

Surface area

A substance which has a larger surface area will evaporate faster as there are more surface molecules which are able to escape.

Temperature of the substance

If the substance is hotter, then evaporation will be faster.

Density

The higher the density, the slower a liquid evaporates.

In the US, the National Weather Service measures the actual rate of evaporation from a standardized "pan" open water surface outdoors, at various locations nationwide. Others do likewise around the world. The US data is collected and compiled into an annual evaporation map. [2] The measurements range from under 30 to over 120 inches (3,000 mm) per year.

Applications

When clothes are hung on a laundry line, even though the ambient temperature is below the boiling point of water, water evaporates. This is accelerated by factors such as low humidity, heat (from the sun), and wind. In a clothes dryer hot air is blown through the clothes, allowing water to evaporate very rapidly.

Combustion vaporization

Fuel droplets vaporize as they receive heat by mixing with the hot gases in the combustion chamber. Heat (energy) can also be received by radiation from any hot refractory wall of the combustion chamber.

Film deposition

Thin films may be deposited by evaporating a substance and condensing it onto a substrate.

See also

http://en.wikipedia.org/wiki/Evaporation

Page 4 of 4

- Atmometer (evaporimeter)
- Crystallisation
- Desalination
- Distillation
- Drying

Jun 05 2009 18:21

- Evaporator
- Evapotranspiration
- Flash evaporation
- Heat of vaporization
- Latent heat
- Pan evaporation
- Transpiration

	То					
From	Solid	Liquid	Gas	Plasma		
Solid	Solid-Solid Transformation	Melting/Fusion	Sublimation	N/A		
Liquid	Freezing	N/A	Boiling/Evaporation	N/A		
Gas	Deposition	Condensation	. N/A	Ionization		
Plasma	N/A	N/A	Recombination/Deionization	N/A		

References

- Sze, Simon Min. Semiconductor Devices: Physics and Technology. ISBN 0-471-33372-7. Has an
 especially detailed discussion of film deposition by evaporation.
- ^ a b Silberberg, Martin A. (2006). Chemistry (4th edition ed.). New York: McGraw-Hill. pp. 431-434. ISBN 0-07-296439-1.
- 2. ^ Geotechnical, Rock and Water Resources Library Grow Resource Evaporation

External links

- MSN Encarta article on evaporation
- Evaporation of water

Retrieved from "http://en.wikipedia.org/wiki/Evaporation"

Categories: Atmospheric thermodynamics | Basic meteorological concepts and phenomena | Materials science | Phase changes | Thin film deposition

Hidden categories: Physics articles needing expert attention | Articles needing expert attention from November 2008

- This page was last modified on 3 June 2009 at 01:00.
- All text is available under the terms of the GNU Free Documentation License. (See Copyrights for details.)

Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a U.S. registered 501(c) (3) tax-deductible nonprofit charity.